

**SIMULTANEOUS ESTIMATION OF AZITHROMYCIN AND  
CEFIXIME IN BULK AND PHARMACEUTICAL FORMULATION BY  
REVERSE PHASE HIGH PERFORMANCE LIQUID  
CHROMATOGRAPHY (RP-HPLC)**

**Usharani Gundala\*<sup>1</sup>, Chandrashekar Bonagiri <sup>2</sup>, Devanna Nayakanti<sup>3</sup>**

<sup>1</sup>Research Scholar, JNTUA, Anantapur, A.P, INDIA.

<sup>2</sup>Professor, MLR Institute of Pharmacy, Hyderabad, A.P, INDIA.

<sup>3</sup>Director of OTRI, JNTUA, Anantapur, A.P, INDIA.

Article Received on  
12 March 2014,

Revised on 05 April 2014,  
Accepted on 28 April 2014

**\*Correspondence for  
Author**

**Usharani Gundala**  
Research Scholar, JNTUA,  
Anantapur, A.P, INDIA.

**ABSTRACT**

Azithromycin and Cefixime are both antibiotics drugs. As there is no UV or HPLC method for the simultaneous estimation of Azithromycin and Cefixime, a need was felt to develop the method for the analysis of both drugs simultaneously. This work concerns with the development and validation of a simple, specific and cost effective RP-HPLC method for simultaneous estimation of Azithromycin and Cefixime in bulk. In addition the developed method was applied to the suitable combined tablet dosage form ie., Zifi-AZ. Chromatography was carried on Kromasil C18 column with mobile phase comprising of

Ortho Phosphoric Acid buffer and Methanol in the ratio of 70:30 v/v. The flow rate was adjusted to 1ml/min with PDA detection at 292 nm. The retention times of Azithromycin and cefixime were found to be 2.8 min, 3.9 min respectively and other replicate standard system suitability parameters are within the limit and uniform. The different analytical parameters such as accuracy, precision, linearity, robustness, limit of detection (LOD), limit of quantification (LOQ) were determined according to the International Conference on Harmonization (ICH) Q2B guidelines. The detector response was linear in the range of 0.5-1.5 mg/ml., 0.4-1.2 mg/ml for Azithromycin and cefixime respectively. The proposed method was successfully applied for the reliable quantification of active pharmaceuticals present in the commercial formulations.


**Keywords:** Azithromycin, Cefixime, HPLC, Methanol, Ortho Phosphoric Acid.

## INTRODUCTION

Azithromycin [9-de-oxy-9a-aza-9a-methyl-9a-homoerythromycin] is an azalide, a subclass of macrolide antibiotics<sup>1</sup>. It acts by inhibiting protein synthesis by binding reversibly to the 'P' site of the 50S ribosomal subunit of the bacteria<sup>2, 3</sup>. It is used for adult and pediatric<sup>4, 5</sup> infections. e.g., respiratory tract infection<sup>1, 6, 7, 8</sup>, skin, soft tissue infections, otitis media<sup>1, 9, 10</sup>, sinusitis, pharyngitis, acute bronchitis, community-acquired pneumonia<sup>11</sup>, cystic fibrosis<sup>11, 12</sup>, tonsillitis<sup>13</sup>, anti-inflammatory in COPD patient<sup>14</sup>, in *P. falciparum* malaria with other antimalarial drugs<sup>15</sup>, typhoid fever<sup>16, 17</sup>.


Cefixime (6R, 7R)-7-[2-(2-amino-4-thiazolyl) glyoxylamido]- 8-oxo-3-vinyl-5-1 – azabicyclo [4.2.0] oct-2- ene-2-carboxylic acid, 7-9z)-[o carboxymethyl]-oxime] is third generation cephalosporin antibiotic. It is under the category of  $\beta$ -lactam antibiotics/cell wall inhibitor. It Acts by inhibiting an enzyme transpeptidase, involved in the building of bacterial cell walls<sup>18</sup>. It is used in lower respiratory tract infections<sup>19, 20, 21</sup>, acute urinary tract infections<sup>21, 22</sup>, biliary tract infections<sup>23</sup>, sinusitis<sup>24</sup>, acute otitis media<sup>25</sup>, peptic ulcer<sup>26</sup>.

**Azithromycin**



**Fig. 1**

**Cefixime**



**Fig. 2**

Combination of Cefixime and Azithromycin has a synergistic effect. The effect of Cefixime against *Neisseria gonorrhoeae* can be significantly enhanced in combination with Azithromycin<sup>27</sup>. This combination is used in treatment of uncomplicated gonococcal urethritis<sup>29</sup>, gonorrhoea<sup>28</sup>, typhoid fever<sup>31, 32</sup>.

Literature survey reveals that HPLC<sup>33</sup>, RP-HPLC<sup>34, 35, 36</sup>, UV-Visible Spectrophotometry<sup>37, 38, 39, 40</sup>, UPLC<sup>41</sup>, methods were reported for the estimation of Azithromycin alone or in combination with other drugs except Azithromycin. As per literature survey, no analytical method has been reported for simultaneous estimation of Cefixime and Azithromycin in pharmaceutical dosage forms. Therefore the present research work, our aim is to develop a

novel, simple, accurate, sensitive, reproducible, economical analytical method to estimate Azithromycin & Cefixime in their combined dosage form in routine analysis.

## **MATERIALS AND METHODS**

### **Chemicals Reagents**

Working standards of pharmaceutical grade Azithromycin & Cefixime were obtained as generous gifts from Dr.Reddy's laboratories (Hyderabad, AP, India) used as such without further purification. The pharmaceutical dosage form used in the study was Zifi-AZ. Methanol (HPLC grade), OPA (AR grade) purchased from Merck specialities Pvt.ltd (Mumbai, India) and double distilled water used for analysis.

### **Instrumentation and chromatographic condition**

Chromatography was carried out on kromasil C18 column with mobile phase comprising of OPA buffer and MeOH in the ratio of 70:30. The flow rate was adjusted to 1ml / min with PDA detection at 292 nm.

### **Preparation of standard solution**

Standard stock solutions of pure drugs were prepared separately by dissolving 10 mg of Azithromycin in 10ml MeOH and 8mg of Cefixime in 10ml MeOH to get concentrations 1mg/ml and 0.8 mg /ml respectively.

### **Preparation of sample solution**

20 tablets were weighed accurately, powdered and equivalent weight was calculated. The equivalent weight of two tablets were taken and dissolved in 100 ml of MeOH to get the concentration 4 mg/ml of cefixime and 5mg /ml of Azithromycin. From stock solution 2 ml was taken and diluted to 10 ml with MeOH to get concentrations 0.8 mg/ml cefixime and 1mg/ml of Azithromycin.

## **RESULTS**

### **Validation**

The developed method was validated with different analytical parameters such as accuracy, precision, linearity, limit of detection, limit of quantification and robustness according to the international conference on harmonization (ICH) Q2B guidelines.

### Precision

Precision of these methods was checked by analyzing the samples at three different time intervals of the same day (intraday precision (table-2)) as well as on different days (interday precision). Robustness for HPLC method was performed by deliberately changing the chromatographic conditions. The flow rate of the mobile phase was changed from 1.0 mL/min to 0.8 mL/min and 1.2 mL/min while ratio of the mobile phase was changed by  $\pm 1\%$ . Results of the Robustness were shown in table-1.

**Table-1: Robustness study**

Parameters	Changes	Retention Time
Azithromycin		
Flow rate (ml/min)	0.8	3.59
	1.2	2.41
Temperature	40°C	2.890
	50°C	2.895
Cefixime		
Flow rate (ml/min)	0.8	4.75
	1.2	3.20
Temperature	40°C	3.84
	50°C	3.86

**Table-2: Precision**

S.N.	Sample Weight	Sample Area-1	Sample Area-2	% Assay	% Assay
1	1418.40	482398	473433	100	100
2	1418.40	481010	471886	100	100
3	1418.40	483932	472590	100	100
4	1418.40	480680	473265	100	100
5	1418.40	480544	473320	100	100
6	1418.40	480109	470543	100	100
<b>Average assay</b>				<b>100</b>	<b>100</b>
<b>STD</b>				<b>0.30</b>	<b>0.24</b>
<b>%RSD</b>				<b>0.30</b>	<b>0.24</b>

### Recovery studies

To check the accuracy of the developed methods and to study the interference of formulation additives, analytical recovery experiments were carried out by standard addition method at 50, 100, 150% levels (table-3 and table-4). From the total amount of drug found percentage recovery was calculated.

**Table-3: Recovery studies of Azithromycin**

S.N.	Spiked Level	Sample Weight (mg)	Sample Area	µg/ml added	µg/ml found	% Recovery
1	50%	709.20	247499	500.00	513.11	103
2	50%	709.20	248653	500.00	515.50	103
3	50%	709.20	246241	500.00	510.50	102
4	50%	709.20	248427	500.00	515.03	103
5	50%	709.20	247861	500.00	513.86	103
6	50%	709.20	241695	500.00	501.07	100
1	100%	1418.40	472870	1000.00	980.34	98
2	100%	1418.40	473206	1000.00	981.03	98
3	100%	1418.40	468669	1000.00	971.63	97
1	150%	2127.60	721023	1500.00	1494.80	100
2	150%	2127.60	722111	1500.00	1497.05	100
3	150%	2127.60	722068	1500.00	1496.96	100
4	150%	2127.60	727747	1500.00	1508.74	101
5	150%	2127.60	726745	1500.00	1506.66	100
6	150%	2127.60	723898	1500.00	1500.76	100

**Table-4: Recovery studies of Cefixime**

S. N.	Spiked Level	Sample Weight (mg)	Sample Area	µg/ml added	µg/ml found	% Recovery
1	50%	709.20	239014	400.00	403.75	101
2	50%	709.20	237237	400.00	400.75	100
3	50%	709.20	235532	400.00	397.87	99
4	50%	709.20	235248	400.00	397.39	99
5	50%	709.20	237349	400.00	400.94	100
6	50%	709.20	233780	400.00	394.91	99
1	100%	1418.40	462329	800.00	780.98	98
2	100%	1418.40	463545	800.00	783.04	98
3	100%	1418.40	461749	800.00	780.00	98
1	150%	2127.60	706867	1200.00	1194.07	100
2	150%	2127.60	705758	1200.00	1192.19	99
3	150%	2127.60	702577	1200.00	1186.82	99
4	150%	2127.60	700858	1200.00	1183.92	99
5	150%	2127.60	700187	1200.00	1182.78	99
6	150%	2127.60	707884	1200.00	1195.79	100

**Linearity****LOD and LOQ**

Limit of Detection (LOD) and Limit of quantification (LOQ) were calculated by using the values of slopes and intercepts of the calibration curves for both the drugs. LOD and LOQ values for Azithromycin were found to be 0.89 and 2.99 and for Cefixime 0.71 and 2.39.

**Robustness**

Method robustness was determined by the small changes in chromatographic conditions like as 0.2ml flow rate and  $\pm 5^\circ\text{C}$  temperature and inject the sample observe the result there were no marked changes compare to other analysis.

## RESULTS AND DISCUSSIONS

Retention times of Azithromycin and Cefixime were found to be 2.8 and 3.9 respectively (as shown in Fig. 3). The detector response was linear in the range of 500-1500 µg/ml, 400-1200 µg/ml for Azithromycin and Cefixime respectively. In the linearity study the regression equation and coefficient of correlation for Azithromycin and Cefixime were found to be ( $y = 4838.1x + 1027.3$ ,  $R^2 = 0.9998$ ), ( $y = 4697.5x + 1638.5$ ,  $R^2 = 0.9996$ ) respectively. Commercial formulations containing Azithromycin and Cefixime were analyzed by the proposed method. A typical chromatogram of marketed formulation is shown in fig. no.3. Six replicate analysis of formulation were carried out and the mean assay values were found close to 100 %. The tailing factors were <2.0 for both the peaks. The elution order was Azithromycin (RT = 2.8 min) and Cefixime (RT = 3.9 min), at a flow rate of 1.0 mL/min. The chromatogram was recorded at 292nm. System suitability was established by injecting standard solution and results are shown in table no.1. The accuracy of the proposed method was determined by recovery studies. It was confirmed from results that the method is highly accurate (table no.3 and 4). Precision (table no.2) was calculated as interday and intraday variations for both the drugs. Percent relative standard deviations for intraday and interday precision for Azithromycin were 0.30 % and 0.36 % and that for Cefixime were 0.24 % and 0.31 % respectively which are well within the acceptable limit of 2 %. For robustness studies in all deliberately varied conditions, the RSD of contents of Azithromycin and Cefixime were found to be well within the acceptable limit of 2%.




Fig. 3




Fig. 4




Fig. 5

## CONCLUSION

The new HPLC method developed and validated for simultaneous estimation of Azithromycin and Cefixime pharmaceutical dosage forms and assured the satisfactory precision and accuracy and also determining lower concentration of each drug in its solid combined dosage form by RP-HPLC method. The method was found to be simple, accurate, precise, economical and rapid and they can be applied for routine analysis in laboratories and is suitable for the quality control of the raw materials, formulations, dissolution studies and can be employed for bioequivalence studies for the same formulation.

## ACKNOWLEDGEMENT

Authors thank full to Rainbow Pharma training lab, Kukatpally, Hyderabad, for providing instruments and analytical support.

**REFERENCES**

1. Ferrara A, Santos CD, Cimbro M: Comparative antimicrobial activity and post-antibiotic effect of azithromycin, clarithromycin and roxithromycin against some respiratory pathogens. *International Journal of Antimicrobial Agents*, 1996; 7:181-186.
2. Sharma HL, Sharma KK “principles of pharmacology” edition 2008: 761-762.
3. Goldman RC, Fesik SW, Doran CC: Role of protonated and neutral forms of macrolides in binding to ribosomes from Grampositive and Gram-negative bacteria. *Antimicrobial Agents Chemotherapy*, 1990; 34:426-431.
4. Treadway G, Reisman A: Tolerability of 3-day, once-daily Azithromycin Dihydrate suspension versus standard treatments for community-acquired paediatric infectious diseases. *International Journal of Antimicrobial Agents*, 2001; 18:427-431.
5. Pacifico L, Chiesa C: Azithromycin Dihydrate in Children: A Critical Review of the Evidence. *Current Therapeutic Research*, 2002; 63:1:54-76.
6. Treadway G, Pontani D, Reisman A: The safety of Azithromycin Dihydrate in the treatment of adults with community-acquired respiratory tract infections. *International Journal of Antimicrobial Agents*, 2002; 19:189-194.
7. Ferrara A, Santos C, Cimbro M, Grassi G: Comparative antimicrobial activity and post-antibiotic effect of Azithromycin Dihydrate, clarithromycin and roxithromycin against some respiratory pathogens. *International Journal of Antimicrobial Agents*, 1996; 7:181-186.
8. Hoepelman I, Mollers M, Schied M: A short (3-day) course of Azithromycin Dihydrate tablets versus a 10-day course of amoxicillin–clavulanic acid (co-amoxiclav) in the treatment of adults with lower respiratory tract infections and effects on long-term outcome. *International Journal of Antimicrobial Agents*, 1998; 9:141-146.
9. Block S, Arrieta A, Seibel M: Single-Dose (30 mg/kg) Azithromycin Dihydrate Compared with 10-Day Amoxicillin/Clavulanate for the Treatment of Uncomplicated Acute Otitis Media: A Double-Blind, Placebo-Controlled, Randomized Clinical Trial. *Current Therapeutic Research*, 2003; 64(A):A30-A42.
10. Rothermel C: Single-Dose Azithromycin Dihydrate for Acute Otitis Media: A Pharmacokinetic/Pharmacodynamic Rationale. *Current Therapeutic Research*, 2003; 64(A): V.
11. Kabra S, Pawaiya R, Lodha R: Long-term daily high and low doses of Azithromycin Dihydrate in children with cystic fibrosis: A randomized controlled trial. *Journal of*



- Cystic Fibrosis, 2010; 9:17-23.
12. Yousef A, Jaffe A: The Role of Azithromycin Dihydrate in Patients with Cystic Fibrosis. *Paediatric Respiratory Reviews*, 2010; 11:108-114.
  13. Baschiera F, Fornai M: Improved Tonsillar Disposition Of Azithromycin Dihydrate Following A 3-Day Oral Treatment With 20 Mg Kg<sup>-1</sup> In Paediatric Patients. *Pharmacological Research*, 2002; 46:1:95-100.
  14. Parnham M: Modulation of neutrophil and inflammation markers in chronic obstructive pulmonary disease by short-term Azithromycin Dihydrate treatment. *European Journal of Pharmacology*, 2005; 517:132-143.
  15. Nakornchai S, Konthiang P: Activity of Azithromycin Dihydrate or erythromycin in combination with antimalarial drugs against multidrug-resistant *Plasmodium falciparum* in vitro. *Acta Tropica*, 2006; 100:185-191.
  16. Ferrera K, Bomasang E: Azithromycin Dihydrate versus First Line Antibiotics in the Therapeutic Management of Documented Cases of Typhoid Fever: A Metaanalysis. *Phil Journal Microbiol Infectious Diseases*, 2004; 33:4:163-168.
  17. Butler T, Sridhar C, Dagac M, Pathak K: Treatment of typhoid fever with Azithromycin Dihydrate versus chloramphenicol in a randomized multicentre trial in India. *Journal of Antimicrobial Chemotherapy*, 1999; 44:2:243-250.
  18. Richard F, Michelle C, Luigi X: *Lippincott's Illustrated Reviews: Pharmacology*, 4th Edition: 360-364.
  19. Quintiliuni R: Cefixime Trihydrate in the Treatment of Patients with Lower Respiratory Tract Infections: Results of US Clinical Trials. *Clinical Therapeutics*, 1996; 18:3:373-390.
  20. Ani R: Comparative, Multicenter Studies of Cefixime Trihydrate and Amoxicillin in the Treatment of Respiratory Tract Infections. *The American Journal of Medicine*, 1988; 85(3A):6-13.
  21. Genvesse I, Carbon C: Cefixime Trihydrate. *International Journal of Antimicrobial Agents*, 1993; 3:1-16.
  22. Iravani A, Richard G: A Double-Blind, Multicenter, Comparative Study of the Safety and Efficacy of Cefixime Trihydrate versus Amoxicillin in the Treatment of Acute Urinary Tract Infections in Adult Patients. *The American Journal of Medicine*, 1988; 85(3A):17-23.
  23. Westphal J, Jehl F, Schloegel M, Monteil H, Brogardl J: Biliary Excretion of Cefixime

- Trihydrate: Assessment in Patients Provided with T-Tube Drainage. *Antimicrobial Agents and Chemotherapy*, 1993; 37:7:1488-1491.
24. Avner S, Buckley J, Pearlman D, Vitanza J: Comparison of Cefixime Trihydrate qd & Amoxicillin tid for acute Sinusitis. *Journal of Allergy and Clinical Immunology*, 1991; 87:1:143.
  25. Johnson C, Susan A, Dennis M: Cefixime Trihydrate compared with amoxicillin for treatment of acute otitis media. *Pediatric Pharmacology and Therapeutics*, 1991:117-122.
  26. Fera M, Carbone M, Foch A: Activity of Cefixime Trihydrate against *Helicobacter pylori*. *International Journal of Antimicrobial Agents*, 1993; 3:105-108.
  27. Verhoef J, Gillissen A: Resistant *Haemophilus influenzae* in community-acquired respiratory tract infections: a role for Cefixime Trihydrate. *International Journal of Antimicrobial Agents*, 2003; 21:501-509.
  28. Furuya R, Nakayama H, Kanayama A: In vitro synergistic effects of double combinations of  $\beta$ -lactams and Azithromycin Dihydrate against clinical isolates of *Neisseria gonorrhoeae*. *Journal of Infection And Chemotherapy*, 2006; 12:4:172-176.
  29. Rahman M, Hoque M, Mamun S, Rahman Z, Sultan Z: Comparison of Single-dose Oral Azithromycin Dihydrate with Azithromycin Dihydrate and Cefixime Trihydrate for treatment of uncomplicated gonococcal urethritis. *International Conference on AIDS*, 2004:11-16.
  30. Ryan K: The *phs* Gene and Hydrogen Sulfide Production by *Salmonella typhimurium*. *Journal of Bacteriology*, 1987; 169:6:2391-2397.
  31. AHFS Drug information. 2009:139-140.
  32. Kucer's use of Antibiotics, 8th edition: 400,804.
  33. Zubata P, Ceresole R, Rosasco R, Pizzorno M: A new HPLC method for Azithromycin Dihydrate quantitation. *Journal of Pharmaceutical and Biomedical Analysis*, 2002; 27:833-836.
  34. Shaikh K, Patil S, Devkhile A: Development and validation of a reversed-phase HPLC method for simultaneous estimation of ambroxol hydrochloride and Azithromycin Dihydrate in tablet dosage form. *Journal of Pharmaceutical and Biomedical Analysis*, 2008; 48:1481-1484.
  35. Singh S, Prakash D, Mani T: Development and Validation of Stability indicating RP-HPLC Method for the estimation of Azithromycin Dihydrate Suspension. *International Journal of ChemTech Research*, 2010; 2:4:1939-1944.

36. Shena Y, Yina C, Sub M, Tua J: Rapid, sensitive and selective liquid chromatography–tandem mass spectrometry (LC–MS/MS) method for the quantification of topically applied Azithromycin Dihydrate in rabbit conjunctiva tissues. *Journal of Pharmaceutical and Biomedical Analysis*, 2010; 52:99-104.
37. Kulikov A, Verushkin A: Development and Validation of a Micellar Liquid Chromatographic Method with UV Detection for Determination of Azithromycin Dihydrate in Tablets and Capsules. *Chromatographia*, 2004; 60:1-2:33-38.
38. Sultana N, Arayne M, Hussain F, Fatima A: Degradation studies of Azithromycin Dihydrate And its Spectrophotometric determination in Pharmaceutical Dosage form. *Pakistan Journal of Pharmaceutical Science*, 2006; 19:2:94-98.
39. Suhagia B, Shah S, Rathod I, Patel H, Doshi K: Determination of Azithromycin Dihydrate in pharmaceutical dosage forms by Spectrophotometric method. *Indian Journal of Pharmaceutical sciences*, 2006; 68:2:242-245.
40. Patel C, Chaudahri C, Patel S, Patel N: Visible spectrophotometric method for estimation of Azithromycin Dihydrate from tablet formulation.
41. Yanamandra R, Chaudhary A, Rao S, Patro B: UPLC Method for Simultaneous Separation and Estimation of Secnidazole, Fluconazole and Azithromycin Dihydrate in Pharmaceutical Dosage Forms. *E-Journal of Chemistry*, 2010; 7:S1:S363-S371.