

STABILITY INDICATING METHOD DEVELOPMENT AND VALIDATION FOR SIMULTANEOUS ESTIMATION OF LOPINAVIR AND RITONAVIR BY USING RP-HPLC

P. S. Raghu*

University College of Pharmaceutical Sciences, Sri Krishnadevaraya University, A.P-515003.

Article Received on
18 Dec. 2017,

Revised on 09 Jan. 2018,
Accepted on 01 Feb. 2018

DOI: 10.20959/wjpr20183-11106

*Corresponding Author

P. S. Raghu

University College of
Pharmaceutical Sciences, Sri
Krishnadevaraya University,
A. P-515003.

ABSTRACT

A simple, Accurate, precise method was developed for the simultaneous estimation of the Lopinavir and Ritonavir in Pharmaceutical dosage form. Chromatogram was run through Inertsil C18 (250mm 4.6mm, 5 μ). Mobile phase containing Orthophosphoric acid buffer and Acetonitrile in the ratio of 44:6 was pumped through column at a flow rate of 1.0 ml/min. Temperature was maintained at 30°C. Optimized wavelength for Lopinavir and Ritonavir was 310nm. Retention time of Lopinavir and Ritonavir were found to be 2.214min and 3.163min. %RSD of the Lopinavir and Ritonavir were and found to be 0.6 and 0.3 respectively. %Recover was Obtained as 99.64% and

99.56% for Lopinavir and Ritonavir. LOD, LOQ values were obtained from regression equations of Lopinavir and Ritonavir were 0.20ppm, 0.61ppm and 0.08ppm, 0.24ppm respectively. Regression equation of Lopinavir $y = 7745x + 22310$, and of Ritonavir is $y = 11440x + 7469$. Retention times were decreased and that run time was also decreased so the method developed was simple and economical that can be adopted in regular Quality control test in Industries.

KEYWORDS: Lopinavir, Ritonavir, RP-HPLC.

INTRODUCTION

Lopinavir¹, is chemically known as (2H) – pyrimidine acetamide N- [[4-(2,6- methyl phenoxy) acetyl]amino]-3-hydroxy 5- phenyl- 1- (phenyl methyl) pentyl, tetrahydro- α - (1- methyl ethyl)- 2- oxo and its empirical formula is C₃₇H₄₈N₄O₅, having a molecular weight of 628.80.




Figure. 1: structure of lopinavir.

Ritonavir², is chemically known as 2,4,7,12- tetra azatridecan- 13oic acid, 10 hydroxy- 2- methyl- 5- (1- methyl ethyl)- 1- [2- (1- methyl ethyl)- 4- thiazoly]- 3,6- dioxo- 8,11- bis(phenyl methyl)- 5- thiazol methyl ester and its empirical formula is C₃₇H₄₈N₆O₅S₂ with a molecular weight of 720.9. Both the drugs were used as antiretroviral agents.




Figure. 2: structure of Ritonavir.

Various analytical methods have been reported for the assay of lopinavir and ritonavir individually or combination with other drugs in biological samples/formulations. They include HPLC³⁻⁶, high performance thin layer chromatography⁷, derivative UV spectrophotometry⁸. Literature survey reveals that no analytical method for determination of lopinavir and ritonavir in combine dosage forms is reported. So it is felt worthwhile to develop a simple, rapid, accurate, precise and more economical high performance liquid chromatographic method for simultaneous estimation of lopinavir and ritonavir in bulk and its combined dosage form.

2. Experimental work

MATERIALS AND INSTRUMENTS

Reference standards of lopinavir and ritonavir were obtained as gift samples from Tenna labs, Hyderabad. Market formulation of this combination Emletra and Ritocom were procured from the local market. HPLC grade acetonitrile and methanol were obtained from Merck (India). Analytical grade potassium hydrogen phosphate buffer and potassium hydroxide

were purchased from SD Fine chemicals, India. Water obtained from Millipore with milli Q system, filtered through 0.45 μ nylon-66 membrane was used for the HPLC work. The LC system consisted of isocratic pump, auto sampler and UV detector. The output signal was monitored and integrated using LC solutions chromatography Manager Software (Prominence HPLC, Shimadzu, Japan).

Mobile phase

A mixture of ortho phosphoric acid phosphate buffer, acetonitrile in the ratio (48:52 v/v) was used as mobile phase which was filtered through a 0.45 μ nylon membrane filter.

Preparation of mixed standard solution of Lopinavir and Ritonavir

About 100 mg of lopinavir and 25 mg of ritonavir were weighed accurately and transferred to 100 ml standard volumetric flask. It was dissolved in mobile phase then the solution was sonicated for about 10 min and the volume was made up to the mark with mobile phase to give a stock solution containing 1mg/ml of lopinavir and 0.25 mg/ml of ritonavir.

Preparation of sample solution of Lopinavir and Ritonavir

To determine the content of the drugs in pharmaceutical formulations, twenty tablets were weighed and pulverized using a mortar and pestle. An amount equivalent to 100 mg of lopinavir and 25 mg of ritonavir, was transferred to a 100ml standard volumetric flask, about 60ml of mobile phase was added and sonicated for about 10minutes. Then volume was made up to the mark with mobile phase and filtered through a 0.45 μ nylon membrane filter. An aliquot portion of the filtrate was further diluted to get final concentration of 500 μ g/ml of lopinavir and 125 μ g/ml of ritonavir. All the determinations were conducted six times to ensure repeatability of the method. The mean peak area of the each drug was calculated.

RESULTS AND DISCUSSION

The purpose of the present study was to develop a rapid and sensitive RP-HPLC method for the simultaneous estimation of lopinavir and ritonavir in combined dosage form using InertsilC18 analytical column with UV detection 310 nm.

Method optimization

To optimize the operating conditions for isocratic RP-LC detection of analytes, a number of parameters such as the mobile phase composition, pH and flow rate were varied. Various ratios (70:30:, 48:52,50:50, 60:40v/v) of buffer: acetonitrile was tested as starting solvent for

system suitability study. The variation in the mobile phase led to considerable changes in the chromatographic parameters like symmetry, capacity factor and retention time. The pH effect showed that optimized conditions are reached when the pH value was 6.0, producing well resolved and sharp peaks for lopinavir and ritonavir. Henceforth, in the present method the ratio of (48:52v/v) potassium hydrogen phosphate buffer: acetonitrile pH adjusted to 6.0 ± 0.1 with 10% sodium hydroxide as a mobile phase, at a flow rate of 1.0 mLmin^{-1} was chosen as optimal conditions. The appropriate wavelength in UV region (310nm) was selected for the measurement of active ingredients in the proposed method. For quantitative determination of lopinavir and ritonavir in formulations, initially mixed standard solution was injected into the column five times and the retention time of lopinavir and ritonavir was found to be 2.24 and 3.163min, respectively (Fig.1).




Figure. 3: chromatogram for lopinavir and ritonavir.

System Suitability

Table. 1: system suitability parameters.

S.No	Name	Retention time	Peak area	Resolution	Tailing factor	USP plate count
1	Lopinavir	2.2	800110	8.3	1.35	7444
2	Ritonavir	3.18	303033		1.44	10664

Validation^[8] The described method has been validated for the simultaneous estimation of lopinavir and ritonavir using following parameters.

Accuracy

Accuracy of the method was demonstrated at three different concentration levels (80-120%) by spiking a known quantity of standard drugs into a analyzed sample in triplicate. The results of accuracy (Table 2) revealed that the method was more accurate.

Precision For the precision of the method, three replicate were injected into the system on two different non consecutive days, in each case %RSD was calculated. Results of precision are given in Table 3, which indicated that the method is precise.

Table. 2: Recovery study of Lopinavir and Ritonavir using the proposed HPLC method.

Sample	Amount added ($\mu\text{g/ml}$)	Recovery (%)	%RSD
Lopinavir	50	99.59	0.4
	100	100.34	0.9
	150	99.00	0.5
Ritonavir	12.5	99.54	0.3
	25	99.30	0.3
	37.5	99.85	0.4

Table. 3: Method precision for Lopinavir and Ritonavir in combined dosage form.

S. No	Drug name	Lablled amount	Amount found in mg	
			Inter day	Intra day
1	Lopinavir	100 mg	99.8 \pm 0.02	99.6 \pm 0.03
2	Ritonavir	25 mg	24.90 \pm 0.03	24.75 \pm 0.08

Linearity To establish linearity of the proposed method, five different sets of drug solution was prepared and analyzed. Standard curves were constructed in the concentration range of 25-150 $\mu\text{g mL}^{-1}$ of lopinavir and 12.5 - 7.5 $\mu\text{g mL}^{-1}$ of ritonavir (Fig.2). Slope, intercept and the correlation coefficient were determined and the regression statistics are shown in Table 3.




Figure. 4: linearity graph of lopinavir.




Figure. 5: linearity curve of Ritonavir.

Table. 4: Analytical Performance Parameters for Lopinavir and Ritonavir.

Parameters	Lopinavir	Ritonavir
Linearity Range	25ppm-150ppm	12.5ppm-75ppm
Regression equation	$y = 7745.x + 22310$	$y = 11440.x + 7469$
Correlation coefficient	0.999	0.999
Slope	7745	11440
Intercept	22310	7469

Limit of detection (LOD) and limit of quantization (LOQ) The limit of detection and limit of quantification for lopinavir and ritonavir were calculated from the linearity data using relative standard deviation of the response and slope of the calibration curve. The limit of detection of a compound is defined as the lowest concentration of analyte that can be detected. The results were tabulated below in table- **Specificity** No interference of peaks were found in the chromatogram indicating that excipients used in the tablet formulation did not interfere with the estimation of the drugs by the proposed method for the simultaneous determination of lopinavir and ritonavir in the combined dosage form, hence the method is specific.

Table. 5: LOD and LOQ.

Drug Name	LOD	LOQ
Lopinavir	0.20	0.61
Ritonavir	0.08	0.24

Robustness In order to demonstrate the robustness of the method, system suitability parameters were verified by making deliberate changes in the chromatographic conditions, viz. change in flow rate by ± 0.05 mL min⁻¹, change in pH of the buffer by ± 0.1 unit and

change in the ratio of mobile phase ($\pm 2\%$ absolute). The method was demonstrated to be robust over an acceptable working range of its HPLC operational parameters.

Table. 6: Robustness.

S. No	Degradation Condition	Lopinavir			Ritonavir		
		% Drug Degraded	Purity Angle	Purity Threshold	% Drug Degraded	Purity Angle	Purity Threshold
1	Acid	4.52	0.206	0.415	4.89	0.427	0.612
2	Alkali	2.99	0.739	0.896	2.94	0.402	0.589
3	Peroxide	1.18	0.214	0.321	1.67	0.492	0.717
4	Thermal	0.91	0.820	1.850	0.75	0.253	0.448
5	UV	0.68	0.766	1.555	0.98	0.372	0.715
6	Water	0.59	0.764	1.548	0.51	0.345	0.687

Table. 7: Summary table for validation parameters.

Validation Parameters	Method developed	
	LOPI	RITO
Range	25-150 $\mu\text{g/ml}$	6.25-37.5 $\mu\text{g/ml}$
Regression Coefficient	0.999	0.999
Slope (m)	7745	11440
Intercept (c)	22310	7469
Regression equation	$Y=7745.x+ 22310$	$y = 11440x + 7469$
Assay	99.54%	99.035
Retention time	2.214	3.18
System Precision (%RSD)	0.8	1.1
Method Precision (%RSD)	0.6	0.3
Accuracy	99.64%	99.56%
LOD	0.20	0.08
LOQ	0.61	0.24
Robustness	0.2%-1.2%	0.7%-1.2%

CONCLUSION

The new stability indicating RP-HPLC method was developed and validated as per guidelines for the simultaneous determination of Lopinavir and Ritonavir in combined pharmaceutical dosage form. The proposed method was found to be accurate, precise, simple, economic, rapid and having good specificity. The developed method can be applied for the assay of commercial tablets containing Lopinavir and Ritonavir in routine quality control analysis.

ACKNOWLEDGEMENT

The author Dr P.S .Raghu gratefully thankful to the Sri Krishnadevaraya University authorities, especially the Honble Vice Chancellor Dr .K Rajagopal, Rector Dr. Lajapathi Rai

and the Registrar Prof.Sudhakar Babu for providing constant encouragement and support during the performance of the research work and preparation of the manuscript.

REFERENCES

1. Hing L. Sham, David A. Betebenner , Xiaoqichen, Ayda saldirar, Sudthida Vasavanonda, Dale J. Kempf, Jacob J.Plattner and Daniel W. “Synthesis and structure activity relationship of a novel series of HIV-1 protease inhibitors encompassing ABT-378 (Lopinavir)”, Bio organic and Medicinal Chemistry Letters, 2002; 12(8): 1185-1187.
2. The Merck Index, An Encyclopedia of Chemicals, Drugs and Biologicals, Merck & Co., Inc., NJ; 2001.
3. Faux. J. Venisse. N, Olivier. J. Bouquet. S. “Rapid high-performance liquid chromatography determination of lopinavir, a novel HIV-1 protease inhibitor, in human plasma”, Chromatographia, 2001; 54: 469-473.
4. Ray John, Pang Edna, Carey Dianne. “Simultaneous determination of indinavir, ritonavir and lopinavir (ABT378) in human plasma by high-performance liquid chromatography”, J.Chromatogr.B. Biomed Sci Appl., 2002; 775: 225-230.
5. S. Frappier, D. Brelih, E. Diarte, B. Ba, D. Ducint; J. L. Pellegrin and M. C. Saux. “Simultaneous determination of ritonavir and saquinavir, two human immunodeficiency virus protease inhibitors, in human serum by high performance liquid chromatography”, J.Chromatogr.B. Biomed Sci Appl., 1998; 714: 384-389.
6. Richard M. W. Hoetelmans, Marjolijn van Essenberg, Monique Profijt, Pieter L. Meenhorst, Jan W. Mulder and Jos H. Beijnen. “High performance liquid chromatographic determination of ritonavir in saliva, cerebro spinal fluid and human plasma”, J.Chromatogr.B. Biomed Sci Appl., 1998; 705(1): 119-126.
7. K.V. Mangoankar, N.D. Prabhu Navelkar, A.V. Sulebhavikar and U.D.Pawar. “HPTLC method for simultaneous determination of lopinavir and ritonavir in capsule dosage form”, E-Journal of Chemistry, 2008; 5(4): 706-712.
8. Carolina Lupi Dias, Ana Maria Bergold, Pedro Edurado Froehlich. “UV-Derivative Spectrophotometric determination of Ritonavir Capsules and Comparison with LC”, Analytical Letters, 2009; 42(12): 1900-1910.